

This is a preview of "BS EN 12823-1:2014". [Click here to purchase the full version from the ANSI store.](#)

BS EN 12823-1:2014



BSI Standards Publication

Foodstuffs — Determination of vitamin A by high performance liquid chromatography

Part 1: Measurement of all-E-retinol and 13-Z-retinol

bsi.

...making excellence a habit.™

This is a preview of "BS EN 12823-1:2014". [Click here to purchase the full version from the ANSI store.](#)

This British Standard is the UK implementation of EN 12823-1:2014. It supersedes BS EN 12823-1:2000 which is withdrawn.

The UK participation in its preparation was entrusted to Technical Committee AW/275, Food analysis - Horizontal methods.

A list of organizations represented on this committee can be obtained on request to its secretary.

This publication does not purport to include all the necessary provisions of a contract. Users are responsible for its correct application.

© The British Standards Institution 2014.
Published by BSI Standards Limited 2014

ISBN 978 0 580 77938 1
ICS 67.050

Compliance with a British Standard cannot confer immunity from legal obligations.

This British Standard was published under the authority of the Standards Policy and Strategy Committee on 31 May 2014.

Amendments/corrigenda issued since publication

| Date | Text affected |
|------|---------------|
|------|---------------|

This is a preview of "BS EN 12823-1:2014". [Click here to purchase the full version from the ANSI store.](#)

EUROPÄISCHE NORM

May 2014

ICS 67.050

Supersedes EN 12823-1:2000

English Version

Foodstuffs - Determination of vitamin A by high performance liquid chromatography - Part 1: Measurement of all-E-retinol and 13-Z-retinol

Produits alimentaires - Détermination de la teneur en vitamine A par chromatographie liquide haute performance
- Partie 1: Dosage du tout-E-rétinol et du 13-Z-rétinol

Lebensmittel - Bestimmung von Vitamin A mit Hochleistungs-Flüssigchromatographie - Teil 1:
Bestimmung von all-E-Retinol und 13-Z-Retinol

This European Standard was approved by CEN on 24 April 2014.

CEN members are bound to comply with the CEN/CENELEC Internal Regulations which stipulate the conditions for giving this European Standard the status of a national standard without any alteration. Up-to-date lists and bibliographical references concerning such national standards may be obtained on application to the CEN-CENELEC Management Centre or to any CEN member.

This European Standard exists in three official versions (English, French, German). A version in any other language made by translation under the responsibility of a CEN member into its own language and notified to the CEN-CENELEC Management Centre has the same status as the official versions.

CEN members are the national standards bodies of Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and United Kingdom.



EUROPEAN COMMITTEE FOR STANDARDIZATION
COMITÉ EUROPÉEN DE NORMALISATION
EUROPÄISCHES KOMITEE FÜR NORMUNG

CEN-CENELEC Management Centre: Avenue Marnix 17, B-1000 Brussels

This is a preview of "BS EN 12823-1:2014". [Click here to purchase the full version from the ANSI store.](#)

| Contents | | Page |
|--|-----------------------------------|-----------|
| Foreword | | 3 |
| 1 | Scope | 4 |
| 2 | Normative references | 4 |
| 3 | Principle | 4 |
| 4 | Reagents | 4 |
| 5 | Apparatus | 7 |
| 6 | Procedure | 8 |
| 7 | Calculation | 10 |
| 8 | Precision | 11 |
| 9 | Test report | 12 |
| Annex A (informative) Examples of HPLC chromatograms | | 13 |
| Annex B (informative) Precision data | | 14 |
| Annex C (informative) Alternative HPLC systems | | 15 |
| Bibliography | | 16 |

This is a preview of "BS EN 12823-1:2014". [Click here to purchase the full version from the ANSI store.](#)

Foreword

This document (EN 12823-1:2014) has been prepared by Technical Committee CEN/TC 275 "Food analysis - Horizontal methods", the secretariat of which is held by DIN.

This European Standard shall be given the status of a national standard, either by publication of an identical text or by endorsement, at the latest by November 2014, and conflicting national standards shall be withdrawn at the latest by November 2014.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. CEN [and/or CENELEC] shall not be held responsible for identifying any or all such patent rights.

This document supersedes EN 12823-1:2000.

This European Standard consists of two parts:

- *Part 1: Measurement of all-E-retinol and 13-Z-retinol;*
- *Part 2: Measurements of β -carotene.*

This European Standard provides the base for the analytical methods. It is intended to serve as a frame in which the analyst can define his own analytical work in accordance to the standard procedure.

According to the CEN-CENELEC Internal Regulations, the national standards organizations of the following countries are bound to implement this European Standard: Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech Republic, Denmark, Estonia, Finland, Former Yugoslav Republic of Macedonia, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, Switzerland, Turkey and the United Kingdom.

This is a preview of "BS EN 12823-1:2014". [Click here to purchase the full version from the ANSI store.](#)

1 Scope

This European Standard specifies a method for the determination of vitamin A in foodstuffs by high performance liquid chromatography (HPLC). This method has been validated in an interlaboratory study with samples of margarine and milk powder with all-E-retinol levels ranging from 653 µg/100 g to 729 µg/100 g and with 13-Z-retinol levels ranging from 30 µg/100 g to 39 µg/100 g. The determination of vitamin A content is carried out by the measurement of all-E-retinol, 13-Z-retinol and β-carotene. This part covers the measurement of all-E-retinol and 13-Z-retinol.

The extract obtained after saponification in this method can be used for the determination of β-carotene, as described in EN 12823-2:2000, *Foodstuffs - Determination of vitamin A by high performance liquid chromatography - Part 2: Measurements of β-carotene*. In this case, the saponification temperature should preferably not exceed 80 °C in order to prevent isomerisation and oxidation of β-carotene.

2 Normative references

The following documents, in whole or in part, are normatively referenced in this document and are indispensable for its application. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

EN ISO 3696, *Water for analytical laboratory use — Specification and test methods (ISO 3696)*

3 Principle

Retinol is saponified by using methanolic or ethanolic potassium hydroxide solution and extracted by an appropriate solvent. The determination is carried out by high performance liquid chromatography (HPLC) with either fluorometric (F) or ultraviolet (UV) detection. The substances are identified on the basis of the retention times and determined by the external standard procedure using peak areas or heights, see [1] to [4].

4 Reagents

During the analysis, unless otherwise stated, use only reagents of recognized analytical grade and water of at least grade 1 according to EN ISO 3696.

4.1 Methanol.

4.2 Ethanol absolute, volume fraction, $\varphi(\text{C}_2\text{H}_5\text{OH}) = 100 \%$.

4.3 Ethanol, $\varphi(\text{C}_2\text{H}_5\text{OH}) = 96 \%$.

4.4 Sodium sulfate, anhydrous.

4.5 KOH solution for saponification, in suitable mass concentrations, e.g. $\rho(\text{KOH}) = 50 \text{ g}/100 \text{ ml}$ or $60 \text{ g}/100 \text{ ml}$, or alcoholic solutions, e.g. 28 g KOH in 100 ml of a mixture of 9 parts per volume of ethanol and 1 part per volume of water.

4.6 Antioxidants, such as ascorbic acid (AA), sodium ascorbate, sodium sulfide (Na_2S), butylated hydroxytoluene (BHT), pyrogallol or hydroquinone.

4.7 Solvents and extraction solvents, such as diethyl ether (peroxide-free), di-isopropylether, light petroleum (boiling range of 40 °C to 60 °C), *n*-hexane, butanol, iso-octane or appropriate mixtures thereof.