

This is a preview of "ISO 13164-3:2013". Click here to purchase the full version from the ANSI store.

First edition  
2013-09-01

---

---

---

## Water quality — Radon-222 — Part 3: Test method using emanometry

*Qualité de l'eau — Radon 222 —  
Partie 3: Méthode d'essai par émanométrie*



Reference number  
ISO 13164-3:2013(E)

This is a preview of "ISO 13164-3:2013". Click here to purchase the full version from the ANSI store.



## **COPYRIGHT PROTECTED DOCUMENT**

© ISO 2013

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

This is a preview of "ISO 13164-3:2013". Click here to purchase the full version from the ANSI store.

## Contents

	Page
<b>Foreword</b>	<b>iv</b>
<b>Introduction</b>	<b>v</b>
<b>1 Scope</b>	<b>1</b>
<b>2 Normative references</b>	<b>1</b>
<b>3 Terms, definitions and symbols</b>	<b>2</b>
3.1 Terms and definitions	2
3.2 Symbols	2
<b>4 Principle</b>	<b>3</b>
<b>5 Sampling</b>	<b>3</b>
5.1 General requirement	3
5.2 Sampling requirement	3
5.3 Sample volume	3
5.4 Container characteristics	3
<b>6 Transportation and storage</b>	<b>4</b>
<b>7 Transfer of radon by degassing</b>	<b>4</b>
7.1 Purpose	4
7.2 Principle	4
<b>8 Detection</b>	<b>4</b>
8.1 Objective	4
8.2 Principle	4
8.3 Silver-activated zinc sulfide ZnS(Ag) scintillation	4
8.4 Air ionization	5
8.5 Semiconductor (alpha-detection)	5
<b>9 Quality assurance and quality control programme</b>	<b>5</b>
9.1 General	5
9.2 Influence quantities	5
9.3 Instrument verification	6
9.4 Method verification	6
9.5 Demonstration of analyst capability	6
<b>10 Expression of results</b>	<b>6</b>
10.1 Activity concentration	6
10.2 Standard uncertainty of the activity concentration	7
10.3 Decision threshold and detection limit	7
10.4 Confidence limits	7
<b>11 Calibration</b>	<b>7</b>
<b>12 Test report</b>	<b>7</b>
<b>Annex A (informative) Examples of measurement methods using scintillation cells</b>	<b>9</b>
<b>Annex B (informative) Example of a measurement method using an ionization chamber</b>	<b>17</b>
<b>Bibliography</b>	<b>23</b>

This is a preview of "ISO 13164-3:2013". Click here to purchase the full version from the ANSI store.

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2, [www.iso.org/directives](http://www.iso.org/directives).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received, [www.iso.org/patents](http://www.iso.org/patents).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

The committee responsible for this document is ISO/TC 147, *Water quality*, Subcommittee SC 3, *Radioactivity measurements*.

ISO 13164 consists of the following parts, under the general title *Water quality — Radon-222*:

- *Part 1: General principles*
- *Part 2: Test method using gamma-ray spectrometry*
- *Part 3: Test method using emanometry*

The following part is under preparation:

- *Part 4: Test method using two-phase liquid scintillation counting*

This is a preview of "ISO 13164-3:2013". Click here to purchase the full version from the ANSI store.

## Introduction

Radioactivity from several naturally occurring and human-made sources is present throughout the environment. Thus, water bodies (surface waters, groundwaters, sea waters) can contain radionuclides of natural and human-made origin.

- Natural radionuclides, including potassium-40, and those of the thorium and uranium decay series, in particular radium-226, radium-228, uranium-234, uranium-238, lead-210, can be found in water for natural reasons (e.g. desorption from the soil and wash-off by rain water) or releases from technological processes involving naturally occurring radioactive materials (e.g. the mining and processing of mineral sands or phosphate fertilizer production and use).
- Human-made radionuclides such as transuranium elements (americium, plutonium, neptunium, curium), tritium, carbon-14, strontium-90 and gamma-emitting radionuclides can also be found in natural waters as they can be authorized to be routinely released into the environment in small quantities in the effluent discharged from nuclear fuel cycle facilities and following their used in unsealed form in medicine or industry. They are also found in water due to the past fallout of the explosion in the atmosphere of nuclear devices and the accidents at Chernobyl and Fukushima.

Drinking-water can thus contain radionuclides at activity concentration which could present a risk to human health. In order to assess the quality of drinking-water (including mineral waters and spring waters) with respect to its radionuclide content and to provide guidance on reducing health risks by taking measures to decrease radionuclide activity concentrations, water resources (groundwater, river, lake, sea, etc.) and drinking water are monitor for their radioactivity content as recommended by the World Health Organization (WHO).

Standard test methods for radon-222 activity concentrations in water samples are needed by test laboratories carrying out such measurements in fulfillment of national authority requirements. Laboratories may have to obtain a specific accreditation for radionuclide measurement in drinking water samples.

The radon activity concentration in surface water is very low, usually below  $1 \text{ Bq l}^{-1}$ . In groundwater, the activity concentration varies from  $1 \text{ Bq l}^{-1}$  up to  $50 \text{ Bq l}^{-1}$  in sedimentary rock aquifers, from  $10 \text{ Bq l}^{-1}$  up to  $300 \text{ Bq l}^{-1}$  in wells, and from  $100 \text{ Bq l}^{-1}$  up to  $1\,000 \text{ Bq l}^{-1}$  in crystalline rocks. The highest activity concentrations are normally measured in rocks with high concentration of uranium (Reference [15]).

High variations in the activity concentrations of radon in aquifers have been observed. Even in a region with relatively uniform rock types, some well water may exhibit radon activity concentration greatly higher than the average value for the same region. Significant seasonal variations have also been recorded (see Annex A).

Water may dissolve chemical substances as it passes from the soil surface to an aquifer or spring waters. The water may pass through or remain for some time in rock, some formations of which may contain a high concentration of natural radionuclides. Under favourable geochemical conditions, the water may selectively dissolve some of these natural radionuclides.

Guidance on radon in drinking-water supplies provided by WHO in 2008 suggests that controls should be implemented if the radon concentration of drinking-water for public water supplies exceeds  $100 \text{ Bq l}^{-1}$ . It also recommended that any new, especially public, drinking-water supply using groundwater should be tested prior to being used for general consumption and that if the radon concentration exceeds  $100 \text{ Bq l}^{-1}$ , treatment of the water source should be undertaken to reduce the radon levels to well below that level (Reference [16]).

This International Standard is one of a series dealing with the measurement of the activity concentration of radionuclides in water samples.

The origin of radon-222 and its short-lived decay products in water and other measurement methods are described generally in ISO 13164-1.