

This is a preview of "ISO 16641:2014". [Click here to purchase the full version from the ANSI store.](#)

First edition  
2014-10-01

---

---

## **Measurement of radioactivity in the environment — Air — Radon 220: Integrated measurement methods for the determination of the average activity concentration using passive solid-state nuclear track detectors**

*Mesurage de la radioactivité dans l'environnement — Air — Radon 220: Méthode de mesure intégrée pour la détermination de l'activité volumique moyenne avec des détecteurs passifs solides de traces nucléaires*



Reference number  
ISO 16641:2014(E)

© ISO 2014



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2014

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
Case postale 56 • CH-1211 Geneva 20  
Tel. + 41 22 749 01 11  
Fax + 41 22 749 09 47  
E-mail [copyright@iso.org](mailto:copyright@iso.org)  
Web [www.iso.org](http://www.iso.org)

Published in Switzerland

This is a preview of "ISO 16641:2014". [Click here to purchase the full version from the ANSI store.](#)

## Contents

	Page
<b>Foreword</b> .....	<b>iv</b>
<b>Introduction</b> .....	<b>v</b>
<b>1 Scope</b> .....	<b>1</b>
<b>2 Normative references</b> .....	<b>1</b>
<b>3 Terms, definitions, and symbols</b> .....	<b>1</b>
3.1 Terms and definitions.....	1
3.2 Symbols.....	3
<b>4 Principle of the measurement method</b> .....	<b>4</b>
<b>5 Equipment</b> .....	<b>5</b>
<b>6 Sampling</b> .....	<b>6</b>
6.1 Sampling objective.....	6
6.2 Sampling characteristics.....	6
6.3 Sampling conditions.....	6
<b>7 Detection method with solid-state nuclear track detectors (SSNTD)</b> .....	<b>7</b>
<b>8 Measurement procedure</b> .....	<b>7</b>
8.1 General.....	7
8.2 Influencing variables.....	8
8.3 Calibration.....	8
<b>9 Expression of results</b> .....	<b>9</b>
9.1 Average thoron activity concentration.....	9
9.2 Standard uncertainty.....	9
9.3 Decision threshold.....	10
9.4 Detection limit.....	10
9.5 Confidence limits.....	11
9.6 Example.....	11
<b>10 Test report</b> .....	<b>12</b>
<b>Annex A (informative) Radon-220 and its decay products</b> .....	<b>14</b>
<b>Bibliography</b> .....	<b>16</b>

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see [www.iso.org/directives](http://www.iso.org/directives)).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see [www.iso.org/patents](http://www.iso.org/patents)).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the WTO principles in the Technical Barriers to Trade (TBT) see the following URL: [Foreword - Supplementary information](#)

The committee responsible for this document is ISO/TC 85, *Nuclear energy, nuclear technologies, and radiological protection*, Subcommittee SC 2, *Radiological protection*.

This is a preview of "ISO 16641:2014". [Click here to purchase the full version from the ANSI store.](#)

## Introduction

Radon isotopes 222, 220, and 219 are radioactive gases produced by the disintegration of radium isotopes 226, 224, and 223, which are decay products of uranium-238, thorium-232, and uranium-235, respectively, are all found in the earth's crust. Solid elements, also radioactive, followed by stable lead are produced by radon disintegration.<sup>[1]</sup>

Radon is considered to be the main source of human exposure to natural radiation. The UNSCEAR (2006) report<sup>[2]</sup> suggests that, at the international level, radon accounts for around 52 % of the global average exposure to natural radiation. Isotope 222 (48 %) is far more significant than isotope 220 (4 %), while isotope 219 is considered negligible.

Recent studies on indoor radon-222 and lung cancer in Europe, North America, and Asia provide strong evidence that radon-222 causes a substantial number of lung cancers in the general population. Current estimates of the proportion of lung cancers attributable to radon-222 range from 3 % to 14 %, depending on the average radon-222 concentration in the country concerned and the calculation methods.<sup>[3]</sup>

Indoor radon-222 concentration is mainly measured by passive detectors that can measure both radon-222 and radon-220 signals.<sup>[4]</sup> If the readings are overestimated, the lung cancer risk is given as a biased estimate when epidemiological studies are carried out. Radon-222 and radon-220 parallel measurements have been carried out in several countries<sup>[4]-[11]</sup> (See [Table A.1](#)). Experiences from field work indicate that there is no correlation among radon-222 and radon-220 and its decay products' concentrations. This implies that one parameter cannot be estimated from the other. Unless radon-220 activity concentration is measured, a correct radon-222 concentration cannot be given with a single use of radon-222 measuring device. Therefore, a specific measurement of radon-220 is justified.

Due to its short half-life, radon-220 disappears very rapidly in the atmosphere. An activity concentration gradient is observed from the walls or grounds to the inner space of the room. Depending on the objective of the measurement (building characteristics, construction material characterization, etc.), the sampling location is to be chosen after taking into account this gradient.

Due to a highest level of radon-222 in air, radon-220 is very difficult to measure alone. This International Standard proposes a measuring method of radon-220 activity concentration using a dual system considering radon-222 and radon-220.

There are many ways of measuring the activity concentration of radon-220 and its decay products. The measuring technique proposed is an integrated measurement method for radon-220 only.