

Second edition  
2019-12

---

---

## Measurement of radioactivity in the environment — Soil —

### Part 5: Strontium 90 — Test method using proportional counting or liquid scintillation counting

*Mesurage de la radioactivité dans l'environnement — Sol —*

*Partie 5: Strontium 90 — Méthode d'essai par comptage  
proportionnel et scintillation liquide*



Reference number  
ISO 18589-5:2019(E)

© ISO 2019



**COPYRIGHT PROTECTED DOCUMENT**

© ISO 2019

All rights reserved. Unless otherwise specified, or required in the context of its implementation, no part of this publication may be reproduced or utilized otherwise in any form or by any means, electronic or mechanical, including photocopying, or posting on the internet or an intranet, without prior written permission. Permission can be requested from either ISO at the address below or ISO's member body in the country of the requester.

ISO copyright office  
CP 401 • Ch. de Blandonnet 8  
CH-1214 Vernier, Geneva  
Phone: +41 22 749 01 11  
Fax: +41 22 749 09 47  
Email: [copyright@iso.org](mailto:copyright@iso.org)  
Website: [www.iso.org](http://www.iso.org)

Published in Switzerland

This is a preview of "ISO 18589-5:2020". Click here to purchase the full version from the ANSI store.

## Contents

	Page
<b>Foreword</b> .....	<b>v</b>
<b>Introduction</b> .....	<b>vi</b>
<b>1 Scope</b> .....	<b>1</b>
<b>2 Normative references</b> .....	<b>1</b>
<b>3 Terms and definitions</b> .....	<b>2</b>
3.1 Terms and definitions.....	2
3.2 Symbols.....	2
<b>4 Principle</b> .....	<b>3</b>
4.1 General.....	3
4.2 Chemical separation.....	3
4.3 Detection.....	4
4.3.1 General.....	4
4.3.2 Source preparation for liquid scintillation counter.....	4
4.3.3 Source preparation for proportional counter.....	4
4.3.4 Background determination.....	4
<b>5 Chemical reagents and equipment</b> .....	<b>5</b>
<b>6 Procedure of strontium desorption</b> .....	<b>5</b>
6.1 Principles.....	5
6.2 Technical resources.....	6
6.2.1 Equipment.....	6
6.2.2 Chemical reagents.....	6
6.3 Procedure.....	6
<b>7 Chemical separation procedure by precipitation</b> .....	<b>7</b>
7.1 Principles.....	7
7.2 Technical resources.....	7
7.2.1 Equipment.....	7
7.2.2 Chemical reagents.....	8
7.3 Procedure.....	8
7.3.1 Separation of alkaline metals and calcium.....	8
7.3.2 Separation of barium, radium and lead.....	9
7.3.3 Separation of fission products and yttrium.....	9
7.3.4 Strontium purification.....	9
7.3.5 Yttrium extraction.....	10
7.3.6 Determination of the chemical yields.....	11
<b>8 Chemical separation procedure by liquid-liquid extraction</b> .....	<b>11</b>
8.1 Principle.....	11
8.2 Technical resources.....	12
8.2.1 Equipment.....	12
8.2.2 Chemical reagents.....	12
8.3 Procedure.....	13
8.3.1 General.....	13
8.3.2 Chemical separation of yttrium.....	13
8.3.3 Source preparation to be measured by PC.....	14
8.3.4 Source preparation to be measured by LSC.....	14
8.3.5 Determination of the chemical yields.....	14
<b>9 Chemical separation procedure by chromatography (crown ether resin)</b> .....	<b>15</b>
9.1 Principles.....	15
9.2 Technical resources.....	15
9.2.1 Equipment.....	15
9.2.2 Chemical reagents.....	15
9.3 Procedure.....	16

This is a preview of "ISO 18589-5:2020". [Click here to purchase the full version from the ANSI store.](#)

9.3.1	General.....	16
9.3.2	Chemical separation of the strontium.....	16
9.3.3	Determination of chemical yield.....	17
<b>10</b>	<b>Measurement.....</b>	<b>17</b>
10.1	General.....	17
10.2	Liquid scintillation counter.....	17
10.3	Gas flow proportional counter.....	17
10.4	Calculation of counting efficiency.....	18
<b>11</b>	<b>Expression of results.....</b>	<b>18</b>
11.1	General.....	18
11.2	Determination of $^{90}\text{Sr}$ in equilibrium with $^{90}\text{Y}$ .....	18
11.2.1	Calculation of the activity per unit of mass.....	18
11.2.2	Standard uncertainty.....	19
11.2.3	Decision threshold.....	19
11.2.4	Detection limit.....	19
11.3	Determination of $^{90}\text{Sr}$ by the $^{90}\text{Y}$ .....	19
11.3.1	Calculation of the activity per unit of mass.....	19
11.3.2	Standard uncertainty.....	20
11.3.3	Decision threshold.....	20
11.3.4	Detection limit.....	21
11.4	Determination of $^{90}\text{Sr}$ in presence of $^{89}\text{Sr}$ when $^{90}\text{Sr}$ is in equilibrium with $^{90}\text{Y}$ .....	21
11.4.1	Calculation of the activity per unit of mass.....	21
11.4.2	Standard uncertainty.....	22
11.4.3	Decision threshold.....	22
11.4.4	Detection limit.....	23
11.5	Confidence limits.....	23
<b>12</b>	<b>Test report.....</b>	<b>23</b>
<b>Annex A (informative) Examples of evaluation models.....</b>		<b>25</b>
<b>Bibliography.....</b>		<b>32</b>

This is a preview of "ISO 18589-5:2020". [Click here to purchase the full version from the ANSI store.](#)

## Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

The procedures used to develop this document and those intended for its further maintenance are described in the ISO/IEC Directives, Part 1. In particular the different approval criteria needed for the different types of ISO documents should be noted. This document was drafted in accordance with the editorial rules of the ISO/IEC Directives, Part 2 (see [www.iso.org/directives](http://www.iso.org/directives)).

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ISO shall not be held responsible for identifying any or all such patent rights. Details of any patent rights identified during the development of the document will be in the Introduction and/or on the ISO list of patent declarations received (see [www.iso.org/patents](http://www.iso.org/patents)).

Any trade name used in this document is information given for the convenience of users and does not constitute an endorsement.

For an explanation on the voluntary nature of standards, the meaning of ISO specific terms and expressions related to conformity assessment, as well as information about ISO's adherence to the World Trade Organization (WTO) principles in the Technical Barriers to Trade (TBT) see the following URL: [www.iso.org/iso/foreword.html](http://www.iso.org/iso/foreword.html).

This document was prepared by Technical Committee ISO/TC 85, *Nuclear energy*, Subcommittee SC 2, *Radiation protection*.

This second edition cancels and replaces the first edition (ISO 18589-5:2009), which has been technically revised.

The main change compared to the previous edition are as follows:

- The introduction has been reviewed accordingly to the generic introduction adopted for the standards published on the radioactivity measurement in the environment.

A list of all parts in the ISO 18589 series can be found on the ISO website.

Any feedback or questions on this document should be directed to the user's national standards body. A complete listing of these bodies can be found at [www.iso.org/members.html](http://www.iso.org/members.html).

## Introduction

Everyone is exposed to natural radiation. The natural sources of radiation are cosmic rays and naturally occurring radioactive substances which exist in the earth and flora and fauna, including the human body. Human activities involving the use of radiation and radioactive substances add to the radiation exposure from this natural exposure. Some of those activities, such as the mining and use of ores containing naturally-occurring radioactive materials (NORM) and the production of energy by burning coal that contains such substances, simply enhance the exposure from natural radiation sources. Nuclear power plants and other nuclear installations use radioactive materials and produce radioactive effluent and waste during operation and decommissioning. The use of radioactive materials in industry, agriculture and research is expanding around the globe.

All these human activities give rise to radiation exposures that are only a small fraction of the global average level of natural exposure. The medical use of radiation is the largest and a growing man-made source of radiation exposure in developed countries. It includes diagnostic radiology, radiotherapy, nuclear medicine and interventional radiology.

Radiation exposure also occurs as a result of occupational activities. It is incurred by workers in industry, medicine and research using radiation or radioactive substances, as well as by passengers and crew during air travel. The average level of occupational exposures is generally below the global average level of natural radiation exposure (see Reference [1]).

As uses of radiation increase, so do the potential health risk and the public's concerns. Thus, all these exposures are regularly assessed in order to:

- improve the understanding of global levels and temporal trends of public and worker exposure;
- evaluate the components of exposure so as to provide a measure of their relative importance;
- identify emerging issues that may warrant more attention and study. While doses to workers are mostly directly measured, doses to the public are usually assessed by indirect methods using the results of radioactivity measurements of waste, effluent and/or environmental samples.

To ensure that the data obtained from radioactivity monitoring programs support their intended use, it is essential that the stakeholders (for example nuclear site operators, regulatory and local authorities) agree on appropriate methods and procedures for obtaining representative samples and for handling, storing, preparing and measuring the test samples. An assessment of the overall measurement uncertainty also needs to be carried out systematically. As reliable, comparable and 'fit for purpose' data are an essential requirement for any public health decision based on radioactivity measurements, international standards of tested and validated radionuclide test methods are an important tool for the production of such measurement results. The application of standards serves also to guarantee comparability of the test results over time and between different testing laboratories. Laboratories apply them to demonstrate their technical competences and to complete proficiency tests successfully during interlaboratory comparisons, two prerequisites for obtaining national accreditation.

Today, over a hundred International Standards are available to testing laboratories for measuring radionuclides in different matrices.

Generic standards help testing laboratories to manage the measurement process by setting out the general requirements and methods to calibrate equipment and validate techniques. These standards underpin specific standards which describe the test methods to be performed by staff, for example, for different types of sample. The specific standards cover test methods for:

- naturally-occurring radionuclides (including  $^{40}\text{K}$ ,  $^3\text{H}$ ,  $^{14}\text{C}$  and those originating from the thorium and uranium decay series, in particular  $^{226}\text{Ra}$ ,  $^{228}\text{Ra}$ ,  $^{234}\text{U}$ ,  $^{238}\text{U}$  and  $^{210}\text{Pb}$ ) which can be found in materials from natural sources or can be released from technological processes involving naturally occurring radioactive materials (e.g. the mining and processing of mineral sands or phosphate fertilizer production and use);

This is a preview of "ISO 18589-5:2020". [Click here to purchase the full version from the ANSI store.](#)

- human-made radionuclides, such as transuranium elements (americium, plutonium, neptunium, and curium),  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{90}\text{Sr}$  and gamma-ray emitting radionuclides found in waste, liquid and gaseous effluent, in environmental matrices (water, air, soil and biota), in food and in animal feed as a result of authorized releases into the environment, fallout from the explosion in the atmosphere of nuclear devices and fallout from accidents, such as those that occurred in Chernobyl and Fukushima.

The fraction of the background dose rate to man from environmental radiation, mainly gamma radiation, is very variable and depends on factors such as the radioactivity of the local rock and soil, the nature of building materials and the construction of buildings in which people live and work.

A reliable determination of the activity concentration of gamma-ray emitting radionuclides in various matrices is necessary to assess the potential human exposure, to verify compliance with radiation protection and environmental protection regulations or to provide guidance on reducing health risks. Gamma-ray emitting radionuclides are also used as tracers in biology, medicine, physics, chemistry, and engineering. Accurate measurement of the activities of the radionuclides is also needed for homeland security and in connection with the Non-Proliferation Treaty (NPT).

This document describes the requirements to quantify the activity of  $^{90}\text{Sr}$  in soil samples after proper sampling, sample handling and test sample preparation in a testing laboratory or in situ.

This document is to be used in the context of a quality assurance management system (ISO/IEC 17025).

This document is published in several parts for use jointly or separately according to needs. These parts are complementary and are addressed to those responsible for determining the radioactivity present in soil, bedrocks and ore (NORM or TENORM). The first two parts are general in nature describe the setting up of programmes and sampling techniques, methods of general processing of samples in the laboratory (ISO 18589-1), the sampling strategy and the soil sampling technique, soil sample handling and preparation (ISO 18589-2). ISO 18589-3 to ISO 18589-5 deal with nuclide-specific test methods to quantify the activity concentration of gamma emitters radionuclides (ISO 18589-3 and ISO 20042), plutonium isotopes (ISO 18589-4) and  $^{90}\text{Sr}$  (ISO 18589-5) of soil samples. ISO 18589-6 deals with non-specific measurements to quantify rapidly gross alpha or gross beta activities and ISO 18589-7 describes in situ measurement of gamma-emitting radionuclides.

The test methods described in ISO 18589-3 to ISO 18589-6 can also be used to measure the radionuclides in sludge, sediment, construction material and products following proper sampling procedure.

This document is one of a set of International Standards on measurement of radioactivity in the environment.

Additional parts can be added to ISO 18589 in the future if the standardization of the measurement of other radionuclides becomes necessary.