

STANDARD

7346-1

Second edition
1996-06-15

Water quality — Determination of the acute lethal toxicity of substances to a freshwater fish [*Brachydanio rerio* Hamilton-Buchanan (Teleostei, Cyprinidae)] —

Part 1:
Static method

*Qualité de l'eau — Détermination de la toxicité aiguë létale de substances vis-à-vis d'un poisson d'eau douce [*Brachydanio rerio* Hamilton-Buchanan (Téléostei, Cyprinidae)] —*

Partie 1: Méthode statique

This material is reproduced from ISO documents under International Organization for Standardization (ISO) Copyright License number IHS/ICC/1996. Not for resale. No part of these ISO documents may be reproduced in any form, electronic retrieval system or otherwise, except as allowed in the copyright law of the country of use, or with the prior written consent of ISO (Case postale 56, 1211 Geneva 20, Switzerland, Fax +41 22 734 10 79), IHS or the ISO Licensor's members.



Reference number
ISO 7346-1:1996(E)

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

International Standard ISO 7346-1 was prepared by Technical Committee ISO/TC 147, *Water quality*, Subcommittee SC 5, *Biological methods*.

This second edition cancels and replaces the first edition (ISO 7346-1:1984), which has been technically revised.

ISO 7346 consists of the following parts, under the general title *Water quality — Determination of the acute lethal toxicity of substances to a freshwater fish [Brachydanio rerio Hamilton-Buchanan (Teleostei, Cyprinidae)]*:

- *Part 1: Static method*
- *Part 2: Semi-static method*
- *Part 3: Flow-through method*

Annexes A, B and C of this part of ISO 7346 are for information only.

© ISO 1996

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the publisher.

International Organization for Standardization
Case Postale 56 • CH-1211 Genève 20 • Switzerland

Printed in Switzerland

This is a preview of "ISO 7346-1:1996". [Click here to purchase the full version from the ANSI store.](#)

Introduction

The three parts of ISO 7346 describe methods of determining the acute lethal toxicity of substances to the zebra fish (*Brachydanio rerio* Hamilton-Buchanan) but it must be emphasized that the recommended use of the zebra fish does not preclude the use of other species. The methodologies presented here may also be used for other species of freshwater, marine or brackish water fish, with appropriate modifications of, for example, dilution water quality and the temperature conditions of the test.

Within the three parts of ISO 7346, a choice can be made between static, semi-static and flow-through methods. The static test, described in this part of ISO 7346, in which the solution is not renewed, has the advantage of requiring simple apparatus, although the substances in the test vessel may become depleted during the course of the test and the general quality of the water may deteriorate. The flow-through method, described in ISO 7346-3, in which the test solution is replenished continuously, overcomes such problems but requires the use of more complex apparatus. In the semi-static procedure, described in ISO 7346-2, the test solutions are renewed every 24 h or 48 h, this method being a compromise between the other two.

The flow-through method can be used for most types of substances, including those unstable in water, but the concentrations of the test substance are determined wherever possible. The static method is limited to the study of substances whose tested concentrations remain relatively constant during the test period. The semi-static method can be used for testing those substances whose concentrations can be maintained satisfactorily throughout the test by renewal of the solutions every 24 h or 48 h. Special arrangements may be necessary for substances which are highly volatile.

To assist in the preparation and maintenance of concentrations of substances which may be lethal at concentrations close to that of their aqueous solubility, a small volume of solvent may be used, as specified in the methods.